



ELSEVIER

Pharmacological Research xxx (2006) xxx–xxx

Pharmacological  
research

www.elsevier.com/locate/yphrs

Review

## Argan oil: Which benefits on cardiovascular diseases?

Mounia Cherki<sup>a,b</sup>, Hicham Berrougui<sup>a</sup>, Anas Drissi<sup>b</sup>,  
Ahmed Adlouni<sup>b</sup>, Abdelouahed Khalil<sup>a,\*</sup>

<sup>a</sup> *Laboratory of Oxidative Stress and Atherosclerosis, Research Centre on Aging, University of Sherbrooke, Sherbrooke, Que., Canada*

<sup>b</sup> *Laboratoire de Recherche sur les Lipoprotéines et L'athérosclérose, Faculté des Sciences Ben M'sik, Casablanca, Morocco*

Accepted 10 February 2006

### Abstract

**Aim:** The argan oil, extracted from argan-tree fruits, has been known for its various pharmacological properties and used as a natural remedy since several centuries. In this review, we present a summary of the results obtained from a survey of the literature on argan oil.

**Data synthesis:** Various studies conducted in vitro or on human and animal models suggest that argan oil could play a beneficial role in cardiovascular diseases prevention and its consumption could protect against atherosclerosis and cancer via a variety of biological mechanisms.

**Conclusion:** Argan oil reduces cardiovascular risk and may be used as anti-atherogenic oil.

© 2006 Elsevier Ltd. All rights reserved.

**Keywords:** Argan oil; Cardiovascular diseases; Blood pressure; Lipids; Oxidation

### Contents

1. Introduction .....	00
2. Argan oil .....	00
3. Biological activity of argan oil .....	00
4. Conclusion .....	00
Acknowledgments .....	00
References .....	00

### 1. Introduction

The link between diet and the development of some pathologies such as arterial hypertension, diabetes, cardiovascular diseases (CVD) and cancer has been the subject of various studies [1,2]. As far as food lipids are concerned, the influence of the ingested quantities and quality in human and animal models was often highlighted [3,4]. Several recommendations based on experimental, epidemiological or nutritional data have shown that the incidence of CVD is positively correlated with saturated fatty acids intake and negatively correlated with unsaturated fatty acids intake, which are abundant but unequally distributed in vegetable oils such as olive, sunflower and linseed oils [3,5].

Indeed, many studies have shown that products rich in components such as tocopherols and phenolic compounds reduce the susceptibility of lipoproteins to lipid peroxidation, which play a crucial role in the development of the atherosclerosis process [6,7]. In addition, virgin olive oil which is rich in oleic acid, decreases the CVD incidence by improving its major risk factors, such as the lipoprotein profile, blood pressure, glucose metabolism and antithrombotic profile [8]. Endothelial function, inflammation and oxidative stress are also positively modulated as consequences of the olive oil consumption [9–11]. Some of these effects are attributed to minor components of virgin olive oil, mainly phenolic compounds [7].

Argan oil, obtained from the pit of *Argania spinosa*, is another vegetable oil rich in oleic acid and linoleic acid with an oleic/linoleic ratio of 1.25. Interestingly, the unsaponifiable fraction of this oil is mainly rich in antioxidant compounds such as sterols, saponin and phenolic compounds, principally

\* Correspondence to: Research Centre on Aging, 1036 Belvedere South, Sherbrooke, Que., Canada J1H 4C4. Tel.: +1 819 829 7131; fax: +1 819 829 7141.

E-mail address: abdelouahed.khalil@usherbrooke.ca (A. Khalil).

$\gamma$ -tocopherol isoform [12]. Considering its rich composition in antioxidant compounds and unsaturated fat, argan oil can be used as a nutritional intervention in the CVD and cancer disease prevention.

## 2. Argan oil

*A. spinosa* (Sapotaceae) is an endemic tree of south-western Morocco where it constitutes the third most common tree. The oleaginous fruits of argan tree furnish edible and marketable oil “argan oil” that provides up to 25% of the dweller daily lipid diet and 9% of annual oil production in Morocco [13]. Argan oil is extracted following multistep procedures from the pit of the fruit by traditional method, press- or solvent-extraction and has been used for nutritional (directly eaten on toasts or used for cooking), cosmetic and medicinal purposes [14]. Recently, it was shown that the geographical origin of the argan fruit and the extraction method used to produce the oil, considerably influence its physicochemical composition and characteristics [15]. Generally, this oil is rich in unsaturated fatty acids (80%) principally oleic and linoleic acids (44.8 and 33.7%, respectively). Interestingly, the unsaponifiable fraction (1% of the oil constituents) of argan oil is mainly rich in antioxidant compounds such as tocopherols, which is present in a higher proportion compared to olive oil (637 mg/kg versus 258 mg/kg, respectively) and especially in its  $\gamma$ -isoform (75%) (Table 1) [12]. Moreover, this non-glyceric fraction is rich in phenolic compounds, principally ferulic and syringic acids (3147 and 37  $\mu$ g/kg, respectively), which are absent in olive oil. Also, it is rich in some sterols such as schottenol (1420 mg/kg) and spinasterol (1150 mg/kg). These two families of sterols, known for their anticancer prop-

erties, are rarely encountered in other vegetable oils [16,17]. Argan oil also contains a non-negligible proportion of squalene, another anti-cancerous product (3140 mg/kg versus 4990 mg/kg in olive oil). These compounds prevent oxidation, contributing to the stability of the oil [18].

## 3. Biological activity of argan oil

The quality of fatty acids contained in argan oil and the abundance of the antioxidant compounds in the unsaponifiable fraction, suggests a putative use for this oil in nutritional prevention against some pathologies such as CVD. However, only few studies have reported the pharmacological effects of this oil and its compounds as anti-hypertensive, hypolipemiant and presenting antioxidant activities in human and animal models [19–23].

It has been known for many years that the unsaturated fatty acid-rich diet could regulate blood pressure, plasma lipids and blood pressure lipoproteins profiles and, consequently, reduce the risk of the development of atherosclerosis and the incidence of CVD [3–5]. This has been exemplified by the beneficial effect of Mediterranean diet attributed to its richness in olive oil [24]. This oil constitutes an important source of monounsaturated fatty acid (MUFA) (73% of oleic acid), while sunflower oil presents a high content in polyunsaturated fatty acids (PUFA) (61.6% of linoleic acid). Interestingly, argan oil presents an equilibrated proportion in MUFA and PUFA (44.8% of oleic acid and 33.7% of linoleic acid). Linoleic acid (C18: 2n – 6) is an essential fatty acid and serves as a precursor for the biosynthesis of arachidonic acid (C20: 4n – 6), which is a precursor of prostaglandin E1 (PGE1), prostacyclin (PGI2) and PGI3 and thromboxane (TXA 2) that are known for their platelet anti-aggregator and vasodilator activities [25]. Kumar and Das [26] suggested that the anti-hypertensive effect of linoleic acid acted by the inhibition of the PUFA on the angiotensin-converting enzyme (ACE) that is known as a powerful vasoactive agent. Also, arachidonic acid has a hypocholesterolemic effect [27]. Indeed, arachidonic-derived fatty acids, particularly  $\gamma$ -linolenic acid, reduced the total cholesterol, VLDL, IDL and LDL-cholesterol concentrations within human and rat sera [28,29]. Those studies pointed out that vegetable oils rich in PUFA decrease the VLDL and LDL-cholesterol and increase the HDL-cholesterol levels in the plasma. Richard et al. [30] also reported that a daily dietary intake of 3–8 mmol  $\gamma$ -linolenic acid reduces serum total cholesterol and LDL-cholesterol levels. It has been suggested that a high-MUFA diet may reduce the risk of CVD by producing LDL particles that are enriched in oleic acid and more resistant to oxidative modifications [4,31]. Hence, this group of fatty acid is not easily oxidizable and is involved in the modulation of the HDL fluidity, which increases their capacity to promote cholesterol efflux [32].

The minor compounds of argan oil, such as plant sterols, may also be implicated in the hypocholesterolemic effect of argan oil [12]. Indeed, the molecular structure of those plant sterols is very similar to that of human cholesterol and thus, plant sterol intake reduces cholesterol absorption by competing with endogenous cholesterol [33,34]. Controlled clinical trials

Table 1  
Chemical composition of argan oil and olive oil [12]

	Argan oil	Olive oil
Tocopherols (mg/kg oil)		
$\gamma$ -tocopherols	480 $\pm$ 7	26 $\pm$ 1
$\alpha$ -tocopherol	35 $\pm$ 1	190 $\pm$ 1
$\delta$ -tocopherol	122 $\pm$ 10	42 $\pm$ 2
Total	637 $\pm$ 18	258 $\pm$ 3
Sterols (mg/100 g oil)		
Schottenol	142 $\pm$ 11	nd
Spinasterol	115 $\pm$ 7	nd
$\delta^{8-22}$ stigmastadiene-3 $\beta$ -ol	9 $\pm$ 1	nd
$\beta$ -sitosterol	nd	156 $\pm$ 3
Campesterol	nd	12 $\pm$ 1
Stigmasterol	nd	nd
Others	29 $\pm$ 1	151 $\pm$ 10
Total	295 $\pm$ 20	319 $\pm$ 14
Phenolic compounds ( $\mu$ g/kg oil)		
Vanillic acid	67 $\pm$ 3	359 $\pm$ 7
Syringic acid	37 $\pm$ 5	0
Ferulic acid	3147 $\pm$ 20	51 $\pm$ 2
Tyrosol	12 $\pm$ 1	19,573 $\pm$ 37
Others	0	773,000 $\pm$ 53
Total	3263 $\pm$ 29	792,983 $\pm$ 99
Squalene (mg/100 g oil)		
	314 $\pm$ 1	499

Data expressed mean  $\pm$  S.E.M.; nd, not detected.

have proven that increasing the amount of sterols in the diet modulates positively the blood lipid profile [35]. Supplementation with 2–3 g of plant sterol (stanol–sterol/day) has been shown to reduce 10–15% of LDL-cholesterol [36]. Berrada et al. [19] and Berrougui et al. [20] have shown an improvement of lipid profile in hyperlipaemia-induced rats by chronic treatment with argan oil. In these two studies, significant decreases in total cholesterol, LDL and body weight were observed, whereas the increased of HDL concentration was not significant [19,20]. These data were also confirmed in human [22,23]. Drissi et al. [22] have demonstrated, in healthy adult subjects living in the south-western of Morocco, that regularly consumption of argan oil is associated with significantly low levels of plasma LDL-cholesterol and lipoprotein(a) compared with healthy subjects living in the same area but are not argan oil consumers. More recently, the first interventional trial on human reported interesting data [23]. In a randomized controlled study, 60 healthy adult men receiving 25 ml/day of argan oil during 3 weeks were compared to a group receiving the same quantity of virgin olive oil. Analyses of the lipid intake showed a reduction in saturated fatty acids with both regimens as compared with the stabilization period. Moreover, serum lipid profile showed a significant increase in HDL-cholesterol and apolipoprotein AI (apo-AI) in both groups. However, LDL-cholesterol and apo-B decreased significantly only in olive oil group as compared with the stabilization period, while triglycerides decreased significantly by 17.5% only in argan oil group. Thus, these results confirm some data obtained previously from rat studies and indicate the triglyceride-lowering effect of argan oil in men.

In the aim to investigate the anti-hypertensive effect of argan oil and its mechanism of action, Berrougui et al. [21] have recently conducted a preventive intervention study of argan oil-rich diet (10 ml/kg) in non-developed hypertensive young SHR rats (4-weeks-old). After 7 weeks of treatment, argan oil administration reduced significantly blood pressure in SHR rats by increasing endothelial response, decreasing thromboxane A2 release, and oxidative stress markers [21]. Thus, these results show the beneficial effect of argan oil in the treatment of hypertension and hyperlipidemia, that is attributed to the PUFA, sterols as well as to other constituents contained in argan oil [21].

In spite of the high content of antioxidants in argan oil, little is known about its action. Recently, some laboratories have investigated in vitro the effect of argan oil minor compounds (tocopherols, sterols and polyphenols) at different concentrations on human-oxidized LDL by incubation with  $\text{CuSO}_4$  [22,37]. LDL lipid peroxidation was evaluated by conjugated diene and malondialdehyde (MDA) formation as well as Vitamin E disappearance. In both studies, results showed that incubation of LDL with tocopherol, sterol and phenolic extracts of argan oil significantly prolonged the lag-phase of LDL peroxidation. Also, the phenolic extracts lowered the rate of lipid peroxidation and reduced the disappearance of Vitamin E in a concentration-dependent manner [22,37]. Incubation of HDL with phenolic extract significantly increased the fluidity of the HDL phospholipid bilayer and the HDL-mediated cholesterol efflux from

THP-1 macrophages. Cherki et al. [38] have also observed an increase (3.7%) of the HDL fluidity of subjects consuming argan oil for 7 weeks without being significant [38].

In parallel, other interesting ex-vivo studies have investigated whether the consumption of argan oil could improve antioxidant status in healthy men [22,39]. The first study showed that a regular consumption of argan oil significantly decreases plasma lipid peroxides (58.3%) and increases significantly the molar ratio of  $\alpha$ -tocopherol/total cholesterol (21.6%) as well as  $\alpha$ -tocopherol concentration (13.4%) when compared to a non-consumer group [22]. In the second study, the results showed that 25 ml/day of argan oil during 3 weeks in healthy men induces a significant increase in the paraoxonase 1 (PON1) activities (29–45%) and Vitamin E concentration (16%) and decrease significantly the lipid peroxides and conjugated diene formation (18 and 8%, respectively) compared to the baseline values. Susceptibility of LDL to lipid peroxidation as well as the maximum diene production (MDP) were significantly reduced (8 and 13%, respectively). These findings confirm the beneficial effect of argan oil on plasma antioxidant status in vitro and show the same effect for argan oil supplementation in men. Thus, argan oil offers an additional natural food supplement to reduce cardiovascular risk.

The protective effect of argan oil is probably due to its high contents of powerful antioxidants, particularly polyphenols, tocopherols and sterols, which are known as powerful antioxidants [10]. These products act by several mechanisms: (1) scavenging of peroxy radicals, which break the peroxidation chain reaction; (2) chelating free  $\text{Cu}^{2+}$  to form redox-inactive complexes and thus reducing metal-catalyzed oxidation of LDL; and (3) inhibiting the binding of  $\text{Cu}^{2+}$  to apolipoproteins and subsequently preventing the modification of amino acid-apo-B protein residue. These hypotheses are also supported by the extended lag-phase and the reduction in the maximum diene production. Thus, LDL may be enriched with different antioxidants of argan oil, which might reduce their susceptibility to lipid peroxidation [40,41]. Gimeno et al. [42] showed that an intake of 25 ml/day of olive oil could increase the resistance of LDL to oxidation. Similarly, several in vitro and in vivo studies in humans as well as in animals have demonstrated that Vitamin E and phenolic compounds extracted from olive oil inhibited oxidation of LDL [6,11,43]. In the case of argan oil, the diminution of LDL susceptibility to lipid peroxidation, could be explained partially by an enrichment with oleic acid that reduce the oxidation potential [44]. In the same time, the marked increase of plasma Vitamin E may be due to abundance of tocopherols in argan oil (636 mg/kg), although the main tocopherol present in this oil is the  $\gamma$ -tocopherol form [12]. The increase of plasma  $\alpha$ -tocopherol levels with argan oil could be consequence of an eventual conversion from  $\gamma$ - to  $\alpha$ -tocopherol because of the close similarity between chemical structures of both molecules [45]. Indeed,  $\gamma$ -tocopherol supplementation simultaneously increases  $\gamma$ - and  $\alpha$ -tocopherol levels. Recent epidemiological, experimental and mechanistic evidences suggest that  $\gamma$ -tocopherol may be a more potent antioxidant and cancer chemopreventive agent than  $\alpha$ -tocopherol [46–48]. For example, it was found that  $\gamma$ -tocopherol have more potent interaction to reactive nitrogen

oxide species than  $\alpha$ -tocopherol [49]. In addition, because Vitamin E is the major natural antioxidant among those present in LDL, it is considered as the first line of defence against lipid peroxidation [6]. This was confirmed by the finding of Nigdikar et al. [50] which showed that the increase in lag-phase with Vitamin E was four- to five-fold greater than with red wine polyphenols.

Also, the antioxidant activity of argan oil could be attributed to its capacity to induce a significant increase in the PON1 activities [39]. Indeed, dietary habits with a high content of vegetable oil could affect paraoxonase and arylesterase activities [51–53]. Nguyen and Sok [54] have demonstrated that the increase of PON1 activities is associated with the unsaturated fatty acid content. Indeed, they showed that monoenoic acids or their phospholipid derivatives play a beneficial role in protecting PON1 from oxidative inactivation as well as in stabilizing PON1 [54]. On the other hand, atherogenic diets have been demonstrated to reduce PON1 activities [51]. Shih et al. [51] showed that C57BL/6J mice fed with a high saturated fat, cholesterol (1.25% (w/w)), and cholate (0.5% (w/w)) diet exhibited reduced serum PON1 activities and hepatic PON1-mRNA levels.

Furthermore, antioxidant substances such as flavonoid quercetin, were shown to reduce the amount of lipoprotein-associated lipid peroxides and preserved PON1 activity [55]. Also, polygonatum extract and Vitamin E reduced oxidative stress and increased serum PON1 activity, but had no effect on PON1-mRNA expression in rabbit fed a high-cholesterol diet [56]. Similarly, Vitamin E supplementation in rats with propylthiouracil-induced hypothyroidism resulted in decreased lipid peroxide levels in plasma and significantly increased serum PON1 activity compared with propylthiouracil alone [57]. The minor polar components of argan oil could also be involved in this change. Thus, increasing plasma PON1 activities potentiate the anti-atherogenic effect of argan oil.

#### 4. Conclusion

This review summarizes the principal findings obtained from argan oil and its beneficial effect as hypolipemiant, hypotensive and antioxidant activities. Indeed, the protective effect of argan oil could be attributed to its interesting chemical composition. It is essentially characterized by the presence of unsaturated fatty acids and antioxidant compounds such as Vitamin E family in which  $\gamma$ -tocopherol account for 70%. With regard to this chemical composition, evidences from observational and experimental studies and interventional trials, suggest that consumption of argan oil may reduce cardiovascular risk by a variety of biological mechanisms including effects on blood pressure, plasma lipid profile and antioxidant status. However, only a few studies have specifically investigated the benefit of a dietary intake of argan oil on CVD. Hence, more investigations are needed to elucidate the bonafide beneficial effects towards CVD as well as its anticancer properties.

#### Acknowledgments

This work was supported by a grant from the Canadian Institute of Health Research (IAO-63150). This support is gratefully

acknowledged. M. Cherki and H. Berrougui are recipients of a studentship from the Quebec-based Strategic Training Program in Aging, (FORMSAV). A. Khalil is Junior 2 fellow of the Fonds de Recherche on Santé du Québec (FRSQ).

#### References

- [1] Shahidi F, Miraliakbari H. Department Omega-3 ( $n-3$ ) fatty acids in health and disease. Part 1. Cardiovascular disease and cancer. *J Med Food* 2004;7:387–401.
- [2] Zhao G, Etherton TD, Martin KR, West SG, Gillies PJ, Kris-Etherton PM. Dietary alpha-linolenic acid reduces inflammatory and lipid cardiovascular risk factors in hypercholesterolemic men and women. *J Nutr* 2004;134:2991–7.
- [3] Trautwein EA, Rieckhoff D, Kunath-Rau A, Erbersdobler HF. Replacing saturated fat with PUFA-rich (sunflower oil) or MUFA-rich (rapeseed, olive and high-oleic sunflower oil) fats resulted in comparable hypocholesterolemic effects in cholesterol-fed hamsters. *Ann Nutr Metab* 1999;43:159–72.
- [4] Nagyova A, Haban P, Klvanova J, Kadrabova J. Effects of dietary extra virgin olive oil on serum lipid resistance to oxidation and fatty acid composition in elderly lipidemic patients. *Bratisl Lek Listy* 2003;104:218–21.
- [5] Keys A, Minotti A, Karvonen MJ. The diet and 15-year death rate in the Seven Countries Study. *Am J Epidemiol* 1986;124:903–15.
- [6] Reaven PD, Khouw A, Beltz WF, Parthasarathy S, Witztum JL. Effect of dietary antioxidant combinations in humans. Protection of LDL by Vitamin E but not by beta-carotene. *Arterioscler Thromb* 1993;13:590–600.
- [7] Mateos R, Dominguez MM, Espartero JL, Cert A. Antioxidant effect of phenolic compounds, alpha-tocopherol, and other minor components in virgin olive oil. *J Agric Food Chem* 2003;51:7170–5.
- [8] Ruiz-Gutierrez V, Muriana FJ, Guerrero A, Cert AM, Villar J. Plasma lipids, erythrocyte membrane lipids and blood pressure of hypertensive women after ingestion of dietary oleic acid from two different sources. *J Hypertens* 1996;14:1483–90.
- [9] Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, et al. Effect of a mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA* 2004;292:1440–6.
- [10] Masella R, Giovannini C, Vari R, Di Benedetto R, Coni E, Volpe R. Effect of dietary virgin olive oil phenols on low density lipoprotein oxidation in hyperlipidemic patients. *Lipids* 2001;36:1195–202.
- [11] Wiseman SA, Mathot JN, de Fouw NJ, Tijburg LB. Dietary non-tocopherol antioxidants present in extra virgin olive oil increase the resistance of low density lipoproteins to oxidation in rabbits. *Atherosclerosis* 1996;120:15–23.
- [12] Khallouki F, Younos C, Soulimani R, Oster T, Charrouf Z, Spiegelhalter B, et al. Consumption of argan oil (Morocco) with its unique profile of fatty acids, tocopherols, squalene, sterols and phenolic compounds should confer valuable cancer chemopreventive effects. *Eur J Cancer Prev* 2003;12:67–75.
- [13] Rahmani M. L'huile d'argan, un produit alimentaire et diététique de qualité. Actes Séminaires sur l'arganier. Division de recherche et d'exploitation forestière. Rabat; 1989, p. 75–82.
- [14] Charrouf Z, Guillaume D. Ethnoeconomical, ethnomedical and phytochemical study of *Argania spinosa* (L.) Skeels. *J Ethnopharmacol* 1999;67:7–14.
- [15] Hilali M, Charrouf Z, Soulhi A, Hachimi L, Guillaume D. Influence of origin and extraction method on argan oil physico-chemical characteristics and composition. *J Agric Food Chem* 2005;53:2081–7.
- [16] Arisawa M, Kinghom D, Cordeii G, Phoebe H, Fansworth R. Plant anticancer agent XXXVI, schottenol glucosid from *Baccharis coridifolia* and *Ipomopsis aggregata*. *Planta Med* 1985;6:544–5.
- [17] Villaseñor IM, Domingo AP. Anticarcinogenicity potential of spinasterol isolated from squash flowers. *Tetragenesis Carcinog Mutagen* 2000;20:99–105.

- [18] Chimi H, Rahmani M, Cillard J, Cillard P. Etude de la fraction phénolique des huiles d'olives vierges et d'argan du Maroc. Actes de l'institut agronomique et vétérinaire 1988;8:17–22.
- [19] Berrada Y, Settaf A, Baddouri K, Cherrah A, Hassar M. Experimental evidence of an antihypertensive and hypocholesterolemic effect of oil of argan, *Argania sideroxylon*. Therapie 2000;55:375–8.
- [20] Berrougui H, Ettaib A, Herrera Gonzalez MD, Alvarez de Sotomayor M, Bennani-Kabchi N, Hmamouchi M. Hypolipidemic and hypocholesterolemic effect of argan oil (*Argania spinosa* L.) in Meriones shawi rats. J Ethnopharmacol 2003;89:15–8.
- [21] Berrougui H, Alvarez de Sotomayor M, Perez-Guerrero C, Ettaib A, Hmamouchi M, Marhuenda E, et al. Argan (*Argania spinosa*) oil lowers blood pressure and improves endothelial dysfunction in spontaneously hypertensive rats. Br J Nutr 2004;92:921–9.
- [22] Drissi A, Girona J, Cherki M, Godas G, Derouiche A, El Messal M, et al. Evidence of hypolipemiant and antioxidant properties of argan oil derived from the argan tree (*Argania spinosa*). Clin Nutr 2004;23:1159–66.
- [23] Derouiche A, Cherki M, Drissi A, Bamou Y, El Messal M, Idrissi-Oudghiri A, et al. Nutritional intervention study with argan oil in man: effects on lipids and apolipoproteins. Ann Nutr Metab 2005;49:196–201.
- [24] Perez-Jimenez F. International conference on the healthy effect of virgin olive oil. Eur J Clin Invest 2005;35:421–4.
- [25] Das UN. Essential fatty acid metabolism in patients with essential hypertension, diabetes mellitus and coronary heart disease. Prostaglandins Leukot Essent Fatty Acids 1995;52:387–91.
- [26] Kumar KV, Das UN. Effect of cis-unsaturated fatty acids, prostaglandins, and free radicals on angiotensin-converting enzyme activity in vitro. Proc Soc Exp Biol Med 1997;214:374–9.
- [27] Hegsted DM, McGrancy RB, Myers ML, Stare FM. Quantitative effects of dietary fat on serum cholesterol in man. Am J Clin Nutr 1965;17:281–95.
- [28] Takada R, Saitoh M, Mori T. Dietary gamma-linolenic acid-enriched oil reduces body fat content and induces liver enzyme activities relating to fatty acid beta-oxidation in rats. J Nutr 1994;124:469–74.
- [29] Fukushima M, Ohhashi T, Ohno S, Saitoh H, Sonoyama K, Shimada K, et al. Effects of diets enriched in  $n-6$  or  $n-3$  fatty acids on cholesterol metabolism in older rats chronically fed a cholesterol-enriched diet. Lipids 2001;36:261–6.
- [30] Richard JL, Martin C, Maille M, Mendy F, Delolanque B, Jacotot B. Effects of dietary intake of gamma-linolenic acid on blood lipids and phospholipid fatty acids in healthy human subjects. J Clin Biochem Nutr 1990;8:75–84.
- [31] Mata P, Varela O, Alonso R, Lahoz C, de Oya M, Badimon L. Monounsaturated and polyunsaturated  $n-6$  fatty acid-enriched diets modify LDL oxidation and decrease human coronary smooth muscle cell DNA synthesis. Arterioscler Thromb Vasc Biol 1997;17:2088–95.
- [32] Sola R, Motta C, Maille M, Bargallo MT, Boisienier C, Richard JL, et al. Dietary monounsaturated fatty acids enhance cholesterol efflux from human fibroblasts. Relation to fluidity, phospholipid fatty acid composition, overall composition, and size of HDL3. Arterioscler Thromb 1993;13:958–66.
- [33] Ostlund RE, Racette SB, Stenson WF. Effects of trace components of dietary fat on cholesterol metabolism: phytosterol, oxysterol and squalene. Nut Rev 2002;60:349–59.
- [34] Plat J, Mensink RP. Effects of plant sterols and stanols on lipid metabolism and cardiovascular risk. Nutr Metab Cardiovasc Dis 2001;11:31–40.
- [35] Plat J, Mensink RP. Plant stanol and sterol esters in the control of blood cholesterol levels: mechanism and safety aspects. Am J Cardiol 2005;96:15D–22D.
- [36] Heinemann T, Axtmann G, von Bergmann K. Comparison of intestinal absorption of cholesterol with different plant sterols in man. Eur J Clin Invest 1993;23:827–31.
- [37] Berrougui H, Cloutier M, Isabelle M, Khalil A. Phenolic-extract from argan oil (*Argania spinosa* L.) inhibits human low-density lipoprotein (LDL) oxidation and enhances cholesterol efflux from human THP-1 macrophages. Atherosclerosis 2006;184:389–96.
- [38] Cherki M, Drissi A, Derouiche A, El Messal M, Bamou Y, Idrissi-Oudghiri A, et al. Beneficial effects of enriched-argan oil diet in healthy men: influence on HDL fluidity and paraoxonase activity. Atherosclerosis 2005;6(Suppl. 1):51.
- [39] Cherki M, Derouiche A, Drissi A, El Messal M, Bamou Y, Idrissi-Oudghiri A, et al. Consumption of argan oil may have an antiatherogenic effect by improving paraoxonase activities and antioxidant status: interventional study in healthy men. Nutr Metab Cardiovasc Dis 2005;15:352–60.
- [40] Fuhram B, Lavy A, Aviram M. Consumption of red wine with meals reduces the susceptibility of human plasma and LDL to lipid peroxidation. Am J Clin Nutr 1995;61:549–54.
- [41] Jialal I, Fuller CJ, Huet BA. The effect of a-tocopherol supplementation on LDL oxidation a dose-response study. Arterioscler Thromb Vasc Biol 1995;15:190–8.
- [42] Gimeno E, Fito M, Lamuela-Raventos RM, Castellote AI, Covas M, Farre M, et al. Effect of ingestion of virgin olive oil on human low-density lipoprotein composition. Eur J Clin Nutr 2002;56:114–20.
- [43] Visioli F, Bellomo G, Montedoro G, Galli C. Low density lipoprotein oxidation is inhibited in vitro by olive oil constituents. Atherosclerosis 1995;117:25–32.
- [44] Zieden B, Kaminskas A, Kristenson M, Olsson AG, Kucinskiene Z. Long chain polyunsaturated fatty acids may account for higher low-density lipoprotein oxidation susceptibility in Lithuanian compared to Swedish men. Scand J Clin Lab Invest 2002;62:307–14.
- [45] Elmadfa I, Kim SW, Reutlinger M, Siewert R. The transformation of gamma-tocopherol to alpha-tocopherol in the animal organism: a generational study in rats. Z Ernahr-Rungswiss 1989;28:36–48.
- [46] Jiang Q, Elsonschwab I, Courtemanche C, Ames BN.  $\gamma$ -tocopherol and its major metabolite, in contrast to  $\alpha$ -tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. Proc Natl Acad Sci U S A 2000;97:11494–9.
- [47] Gao R, Stone WL, Huang T, Papas AM, Qui M. The uptake of tocopherols by RAW264.7 macrophages. Nutr J 2002;1:2.
- [48] Huang HY, Alberg AJ, Norkus E, Hoffman SC, Comstock GW, Helzlsouer KJ. Prospective study of antioxidant micronutrients in the blood and the risk of developing prostate cancer. Am J Epidemiol 2003;157:335–44.
- [49] Cooney RV, Franke AA, Harwood PJ, Hatch-Pigott V, Custer LJ, Mordan LJ. Gamma-tocopherol detoxification of nitrogen dioxide: superiority to alpha-tocopherol. Proc Natl Acad Sci U S A 1993;90:1771–5.
- [50] Nigdikar SV, Williams NR, Griffin BA, Howard AN. Consumption of red wine polyphenols reduces the susceptibility of low-density lipoproteins to oxidation in vivo. Am J Clin Nutr 1998;68:258–65.
- [51] Shih DM, Gu L, Hama S. Genetic-dietary regulation of serum paraoxonase expression and its role in atherogenesis in a mouse model. J Clin Invest 1996;97:1630–9.
- [52] Durrington PN, Mackness B, Mackness MI. The hunt for nutritional and pharmacological modulators of paraoxonase. Arterioscler Thromb Vasc Biol 2002;22:1248–50.
- [53] Aviram M, Rosenblat M. Paraoxonases and cardiovascular diseases: pharmacological and nutritional influences. Curr Opin Lipidol 2005;16:393–9.
- [54] Nguyen SD, Sok DE. Beneficial effect of oleoylated lipids on paraoxonase. I: protection against oxidative inactivation and stabilization. Biochem J 2003;375:275–85.
- [55] Mackness M, Mackness B, Durrington P, Connelly P, Hegele R. Paraoxonase: biochemistry, genetics and relationship to serum lipoproteins. Curr Opin Lipidol 1996;7:69–76.
- [56] Jeon SM, Lee MK, Park YB, Park HM, Choi MS. Polygonatum rhizoma affects antioxidant defense systems without changing mRNA expression in diet-induced hypercholesterolemic rabbits. J Med Food 2004;7:358–65.
- [57] Sarandol E, Tas S, Dirican M, Serdar Z. Oxidative stress and serum paraoxonase activity in experimental hypothyroidism: effect of Vitamin E supplementation. Cell Biochem Funct 2005;23:1–8.